arXiv:cond-mat/0609050v1 [cond-mat.other] 4 Sep 2006

Quantum phase transitions of light.

Andrew D. Greentree,[1,*] Charles Tahan,[2,1] Jared H. Cole,[1] and L. C. L. Hollenberg[1]

[1] *Centre for Quantum Computer Technology, School of Physics, The University of Melbourne, Victoria 3010, Australia.*

[2] *Cavendish Laboratory, University of Cambridge, JJ Thomson Ave, Cambridge CB3 0HE, United Kingdom*

(Dated: February 6, 2008)

Recently, condensed matter and atomic experiments have reached a length-scale and temperature regime where new quantum collective phenomena emerge. Finding such physics in systems of photons, however, is problematic, as photons typically do not interact with each other and can be created or destroyed at will. Here, we introduce a physical system of photons that exhibits strongly correlated dynamics on a meso-scale. By adding photons to a two-dimensional array of coupled optical cavities each containing a single two-level atom in the photon-blockade regime, we form dressed states, or polaritons, that are both long-lived and strongly interacting. Our zero temperature results predict that this photonic system will undergo a characteristic Mott insulator (excitations localised on each site) to superfluid (excitations delocalised across the lattice) quantum phase transition. Each cavity's impressive photon out-coupling potential may lead to actual devices based on these quantum many-body effects, as well as observable, tunable quantum simulators.

The Jaynes-Cummings [1] model is arguably the most important model for understanding light-matter interactions. It describes the interaction of a single, quasi-resonant optical cavity field with a two-level atom. The coupling between the atom and the photons leads to optical nonlinearities and an effective photon-photon repulsion. Perhaps the most extreme demonstration of this photonic repulsion is photon blockade, demonstrated recently by Birnbaum *et al.* [2], where photonic repulsion prevents more than one photon from being in the cavity at any one time. Photon blockade was initially theoretically described with a four-state system [3], with multiplication of the weak Kerr nonlinearity effected by placing a large number of atoms within each cavity. However, it was quickly realised that the photonic blockade mechanism does not persist in the limit of many atoms [4], rapidly degrading as the number of atoms per cavity is increased [5]. Later Rebic *et al.* showed that the nonlinear interaction afforded by placing a single two-level atom inside a cavity would suffice for realising photon blockade [6]. This observation was highly significant as it allowed the full weight of the Jaynes-Cummings model to be used to attack and understand this problem.

To create an atom-photon system whose dynamics mirror those traditionally associated with strongly interacting condensed matter systems, we consider a two-dimensional array of photonic bandgap cavities. Each cavity contains a single two-level atom, quasi-resonant with the cavity mode. Evanescent coupling between the cavities due to their proximity allows inter-cavity photon hopping. This configuration is depicted schematically in Fig. 1(a), where we have explicitly chosen three nearest neighbours per cavity (coordination number $z = 3$), for reasons explained below. Because we are considering small cavities, with volumes of order λ^3 where λ is the wavelength of the light, there will be strong atom-photon couplings that will dominate over the spontaneous emission. Fig. 1(b) shows the pertinent energy scales within one of the cavities. In the classical limit without two-level atoms, an array of coupled photonic bandgap cavities has been described for novel waveguiding applications [7, 8], and in the quantum regime a two cavity arrangement has been proposed as a Q-switch [9]. We consider here for the first time, the rich dynamics of a two dimensional lattice of *quantum* cavities.

Interest in interacting boson systems grew out of work on the metal-insulator transition in Fermi systems [10]. The seminal paper of Fisher *et al.* [11] employed an infinite range mean-field theory to qualitatively describe the superfluid-insulator phase transitions of a bosonic gas in random and periodic potentials, with and without disorder. Recently, it has been shown that this Bose-Hubbard model can be realized in cold atom optical lattices [12]. The Bose-Hubbard Hamiltonian seeks to describe the many-body dynamics of strongly interacting bosons,

$$\mathcal{H}^{BH} = \sum_{i,j} t_{ij} a_j^\dagger a_i + U \sum_i a_i^\dagger a_i^\dagger a_i a_i, \qquad (1)$$

where i, j range over all sites in the lattice (fixed in our case), t_{ij} is the hopping energy of bosons between sites i and j, U is the on-site repulsion between particles, and a_i is the bosonic annihilation operator on site i. In our photonic-atom superlattice, the Jaynes-Cummings interaction provides the effective on-site repulsion, and the hopping term is represented by the evanescent coupling between the cavities. However, our physical system is fundamentally different from the Bose-Hubbard case: in particular, the effective U is not constant in the Jaynes-Cummings system, but decreases as the number of pho-

tons in each cavity goes up; and the conserved particles here are not pure bosons, but dressed photons or polaritons, which are a mixture of the spinor atom and photons in each cavity. Thus, the hopping term corresponds to the overlap of these extended composite photon-atom states, which are the conserved particles in the model, *not* the photons.

To motivate the search for Hubbard model type interactions within a superlattice of photonic bandgap cavities, we first discuss the Hamiltonian for a single two-level atom in a quasi-resonant cavity,

$$\mathcal{H}^{JC} = \epsilon\sigma_+\sigma_- + \omega a^\dagger a + \beta\left(\sigma_+ a + \sigma_- a^\dagger\right), \qquad (2)$$

where σ_+ and σ_- (a, $a^\dagger$) correspond to the atomic (photonic) raising and lowering operators, respectively. The transition energy of the atomic system is ϵ, the cavity resonance is ω and the cavity mediated atom-photon coupling is β, which is implicitly assumed to be real for our purposes. Defining the atomic states as $|g\rangle$ and $|e\rangle$ for ground and excited state respectively, we introduce the detuning, $\Delta \equiv \omega - \epsilon$. The eigenstates of Eq. 2 are the dressed states [13], which we label as $|\pm, n\rangle$ defined in the Methods section, where n is the number of excitations in the cavity. The ground state for the Jaynes-Cummings Hamiltonian is qualitatively different from the other dressed states, implying a non-trivial form of the raising operator [14]. This constitutes a further departure from usual Hubbard-like condensed-matter models, where the raising operator is not dependent on the number of excitations.

In Fig. 2 we show some of the eigenvalues of the Jaynes-Cummings system with the photon energy subtracted for ease of comparison, i.e. we are plotting $E_{|\pm,n\rangle} - n\omega$. The eigenenergies are well known and are $E_{|\pm,n\rangle} = n\omega \pm \chi(n) - \Delta/2$ where $\chi(n) = \sqrt{n\beta^2 + \Delta^2/4}$ is the n photon generalised Rabi frequency. Displaying the eigenvalues in this way immediately allows us to connect the Jaynes-Cummings Hamiltonian with the on-site repulsion in the Hubbard Hamiltonian. The on-site repulsion is evinced by the increasing energy separation with n.

The Hamiltonian for our extended Hubbard-like system is given by a combination of the Jaynes-Cummings Hamiltonian with photon hopping between cavities and the chemical potential term,

$$\mathcal{H} = \sum_i \mathcal{H}_i^{JC} + \sum_{\langle i,j\rangle} \kappa_{ij}\left(a_i^\dagger a_j + a_j^\dagger a_i\right) - \sum_i \mu_i N_i, \quad (3)$$

where the inter-cavity hopping occurs with frequency $\kappa_{ij} = \kappa$ for nearest neighbours, and $\kappa = 0$ otherwise, N_i is the total number of atomic and photonic excitations (the conserved quantity in our system) and μ_i is the chemical potential at site i in the grand canonical ensemble. For our proof of concept calculation we assume zero disorder and $\mu_i = \mu$ for all sites. To most effectively explore the important regime where on-site repulsion dominates over hopping, we consider a lattice with as few nearest neighbours as possible to achieve a two-dimensional network, that is three nearest neighbours. Altering the number of nearest neighbours does not qualitatively affect our results.

To gain insight over the properties of the full Hamiltonian, we will employ a mean-field approximation. Mean-field theories [15, 16, 17] give good qualitative and quantitative descriptions of these systems, comparing well to Monte Carlo simulations [18, 19]. These mean-field approaches have also been extended to dipolar bosons [20] and boson-fermion atomic mixtures [21, 22], as well as to the theory of exciton and exciton-polariton (electron-hole pair plus photon) condensation [23]. We introduce a super-fluid order parameter, $\psi = \langle a_i\rangle$, and employ the decoupling approximation $a_i^\dagger a_j = \langle a_i^\dagger\rangle a_j + \langle a_j\rangle a_i^\dagger - \langle a_i^\dagger\rangle\langle a_j\rangle$. The resulting mean-field Hamiltonian can be written as a sum over single sites,

$$\begin{aligned}\mathcal{H}^{MF} \;=\; \sum_i \Big\{ & \mathcal{H}_i^{JC} - z\kappa\psi\left(a_i^\dagger + a_i\right) + z\kappa\,|\psi|^2 \\ & -\mu\left(a_i^\dagger a_i + \sigma_i^+\sigma_i^-\right)\Big\}, \end{aligned} \qquad (4)$$

where $z = 3$ is the number of nearest neighbours. To obtain the system's zero temperature properties, we use the procedure of Refs. [16, 17] which is outlined in the Methods section. When $\psi = 0$ we have a Mott phase, characterised by a fixed number of excitations per site with no fluctuations, and $\psi \neq 0$ indicates a superfluid phase. The boundary between the $\psi = 0$ and $\psi \neq 0$ phases denotes where a quantum phase transition in this system will occur: *a quantum phase transition of light.* In general, we expect that when photon-photon coupling (on-site repulsion) dominates over hopping, the system should be in a Mott phase, and when the converse is true, the system will be in a superfluid phase.

We can determine the extent of the Mott lobes in the limit of very small hopping from inspection of Eqs. 2 and 4. First note that with our definition, $E_{|-,n\rangle} < E_{|+,n\rangle}$, we only need consider the negative branch for the purposes of determining the ground state. Furthermore, a change in the total number of excitations per site will occur when $E_{|-,n+1\rangle} - \mu(n+1) = E_{|-,n\rangle} - \mu n$. We can determine the critical chemical potential, $\mu_c(n)$, where the system will change from n to $n+1$ excitations per site as

$$\mu_c(n) = \omega - \left[\chi(n+1) - \chi(n)\right]. \qquad (5)$$

These boundary chemical potentials are shown in Fig. 3 as a function of detuning in the limit of very small κ, and these also account for the asymptotes in Fig. 4. In Fig. 4 we show the complete phase diagram of the mean field solution. Because of the large parameter space, there are several cases that we consider. Note that our results are only shown for $\mu < \omega$ as in the limit $\mu \gtrsim \omega$ the minimisation method does not converge [$\mu_c(\infty) = \omega$],

although in this limit, the quantum ground state should correspond to a coherent state of excitations (superfluid phase).

The dynamics illustrated in Fig. 4 are extremely rich. Hubbard-like dynamics can be seen in the three subfigures of Fig. 4, which show ψ as a function of κ and μ for three different detunings. In this plot, $\psi = 0$ corresponds to stable Mott lobes, with the number of excitations increasing with μ. The regions to the right correspond to $\mu \neq 0$, and in these regions the system will be found in a superfluid phase, i.e. the stable ground state at each site corresponds to a coherent state of excitations over the $|-, n\rangle$ branch. The size of the Mott lobes varies with Δ, with the largest Mott lobes found on resonance.

To confirm the number of photons in each Mott lobe, we show in Fig. 5 the average number of excitations per site in the grand-canonical ensemble, $\rho = -\partial E_g(\psi = \psi_{min})/\partial\mu$, as a function of κ and μ for $\Delta = 0$. These plataues indicate regions of constant density and incompresibility, both characteristic of the gapped Mott insulator phase.

POTENTIAL IMPLEMENTATIONS

Until now, we have deliberately avoided specific implementations, as the above analysis should be applicable to all atom-photon systems with strong coupling. In this section we specifically consider some candidate implementations that may be realisable with present-day or near-term technology.

One candidate system would be a photonic bandgap structure in diamond, with individual negatively-charged nitrogen-vacancy (NV) centres placed in each cavity. Micromachining of diamond is presently being explored for similar purposes, see [24, 25], and modelling has shown that suitable cavities are in principle fabricatable [26]. The application of diamond for quantum computing and quantum optics applications has recently been reviewed [27]. For NV diamond, the two-state transition of interest is at 637 nm $\sim 3 \times 10^{15}$ Hz. Calculations presented in Ref. 9 showed that λ^3 single-mode photonic bandgap cavities would have $\beta \sim 10^{10}$Hz. Assuming that photon hopping limits the cavity Q, we can approximate tunneling frequency by $\kappa = \omega/Q$, so for $\kappa \leq \beta$ would require $Q \geq 10^5$. This is demanding, but we note that $Q \sim 10^7$ has been achieved in silicon-on-silica photonic-bandgap cavities [28], although we stress that complete modeling is necessary to precisely determine the required geometry. If the diamond substrate is ultra-purity Type IIa (low nitrogen) diamond, individual NV centres can be implanted using single-ion implantation techniques [27, 29]. Finally we note that the resonance frequencies for the photonic bandgap cavities will be extremely difficult to tune post-creation, however the Stark effect can be used to tune the NV centres as required [30] to allow an exploration (either statically or dynamically) of the phase space shown in Fig. 4.

Another candidate system where such effects may be observable is the microwave strip line resonator for circuit QED recently demonstrated by Walraff *et al.* [31]. In such systems, the effective mode volume of the cavity can be much less than λ^3, with $10^{-6}\lambda^3$ demonstrated in Ref. 31, with the rate of loss of photons out of the cavity (analogous to the κ described in this paper) being $\sim 10^3$ Hz. When such values are combined with their atom-photon couplings, which were of order $\beta \sim 10^7$ Hz, then the realised cavities would seem to be well-suited to observing the effects we have described. One advantage of the strip lines over the photonic bandgap case, however, would be that generating structures with controlled (and perhaps varying) numbers of nearest neighbours would be much easier.

The above two examples do not constitute the limit of possible implementations. It may be possible to achieve these phase transitions using Rydberg atoms in high-Q superconducting cavities. Such systems have shown impressive results including generation of one and two-photon states in the cavity [32]. An alternative architecture would be an array of microcavities containing single atoms, with the cavities connected by optical fibres [33]. Again, this system would appear to offer the desirable properties of high atom-photon coupling, with relatively low hopping terms, and classical routing would permit interesting lattice structures. Optical quantum dots, for example InAs structural quantum dots in a photonic bandgap lattice, may also afford interesting possibilities for the realisation of these (or similar) quantum phase transitions.

CONCLUSIONS

We have shown how a Mott insulator to superfluid transition can be realized with atom-dressed photons in a cavity superlattice. Calculations for zero temperature give a phase diagram analogous to, but fundamentally different from, the Bose-Hubbard model.

If one could Q-switch the cavities, for example by bringing each cavity into close proximity to a near field probe, it may be possible to out-couple all of the cavities simultaneously. This would generate a self-ordered, two-dimensional array of single photon emitters, which could be extremely beneficial for quantum information applications. Because there is significant flexibility in constructing exotic lattice geometries, one may be able to engineer an extended Bose-Hubbard system suitable for topological quantum computing [34] where fault-tolerance is a natural consequence. Alternatively, simulation of quantum systems by regime replication (where the Hamiltonian of the simulator system is transformed to be equivalent to the Hamiltonian of some unknown system) is an

attractive application.

Similar though modified dynamics found in other Hubbard-like systems should also be realizable in the atom-dressed photon superlattice, such as a glass phase [11], stripes and other symmetry breaking phases [35], super-solid behavior [36], etc. More importantly, since the excitation spectrum of a fluctuation is gapped in the Mott-insulator region, fluctuations due to temperature are exponentially suppressed. Thus, one will be able to observe a phase that is formally not a Mott insulator, but experimentally has very similar features [11, 16]. Disorder-induced destruction of the Mott state [11] is also suppressed by this excitation-hole gap, and system tunability as via the Stark shift may allow for manual correction of these irregularities. Because of the Mott phase's robustness, devices based on this effect at non-zero temperature should be possible in this system. We also note that signatures of quantum many-body phenomena should appear in 1D or finite arrays of cavities.

Note Added: Whilst we were preparing our manuscript, we became aware of two related works, one by Hartmann *et al.* [37], and one by Angelakis *et al.* [38], treating arrays of coupled quantum cavities. The former [37] considers a four-state scheme for realising the required quantum nonlinearity with time domain analysis. The latter [38] performs a similar analysis explicitly considering two-state systems per cavity in a 1-D arrangement to realise an XY chain. Neither paper demonstrates Mott lobes as done here, nor did they perform a mean field analysis. As such, both works are qualitatively different from the present one, but demonstrate that photonic systems are ripe for the exploration of condensed matter physical effects.

METHODS

The eigenvalues and eigenvectors of the Hamiltonian of Eq. 2 are

$$|\pm, n\rangle = \frac{\left[-\frac{\Delta}{2} \pm \chi(n)\right]|g, n\rangle + \beta\sqrt{n}|e, n-1\rangle}{\sqrt{2\chi^2(n) \pm \chi(n)\Delta}} \;\forall n \geq 1, \tag{6}$$

with eigenvalues

$$E_{|\pm,n\rangle} \;=\; n\omega \pm \chi(n) - \Delta/2. \tag{7}$$

Because $n \geq 1$, the above definition for the dressed states does not extend to $n = 0$, and so we must define the ground state for the dressed state system as $|g, 0\rangle$, with $E_{|g,0\rangle} = 0$.

To calculate the ground state wavefunction of our extended Jaynes-Cummings Hamiltonian we must calculate the ground-state energy E_g as a function of ψ as follows. First: obtain the matrix elements of H_i^{MF} in the onsite basis $\{|g, 0\rangle, |e, 0\rangle, |g, 0\rangle, |e, 1\rangle, \ldots\}$ truncated at a finite value of n states, n_{max}. Second: diagonalize this matrix (in the truncated basis) and identify the lowest eigenvalue, E_g. Third: increase n_{max} until E_g converges to its value at $n_{max} = \infty$. Finally: minimize E_g with respect to ψ for different values of κ, ω, and Δ to obtain the phase diagram. At $T = 0$ this approach is operationally equivalent [17] to the Gutzwiller Ansatz (GA) variational wave function technique [39, 40] often used, but can be extended to finite temperatures.

For clarity, we will write out the first few terms of the mean-field Hamiltonian in the onsite basis, $|g, 0\rangle$, $|e, 0\rangle$, $|g, 1\rangle$, etc.

$$\mathcal{H}^{MF} = \begin{bmatrix} \boxed{0} & 0 & -z\kappa\psi & & & \\ 0 & \boxed{\epsilon-\mu} & \boxed{\beta} & -z\kappa\psi & 0 & \\ -z\kappa\psi & \boxed{\beta} & \boxed{\omega-\mu} & 0 & -\sqrt{2}z\kappa\psi & \\ & -z\kappa\psi & 0 & \boxed{\epsilon+\omega-2\mu} & \boxed{\sqrt{2}\beta} & -z\sqrt{2}\kappa\psi \\ & 0 & -\sqrt{2}z\kappa\psi & \boxed{\sqrt{2}\beta} & \boxed{2\omega-2\mu} & 0 \\ & & & -z\sqrt{2}\kappa\psi & 0 & \boxed{\epsilon+2\omega-3\mu} \\ & & & 0 & -\sqrt{3}z\kappa\psi & \boxed{\sqrt{3}\beta} \\ & & & & & \ddots \end{bmatrix}, \tag{8}$$

where the entries in the boxes correspond to the usual block diagonal form of the Jaynes-Cummings Hamiltonian with the addition of the superfluid order parameter, and the number of excitations in each block increases by one as we move diagonally from block to block from the ground state $|g, 0\rangle$. The critical chemical potentials can be immediately determined by equating the lower eigenstates (negative dressed-state branch) from neighbouring blocks, with $\kappa = 0$, which gives the result in Eq. 5.

ACKNOWLEDGEMENTS

The authors would like to acknowledge useful discussions with P. Eastham, M. Friesen, D. Jamieson, R. Joynt, R. Kalish, P. Littlewood, A. Martin, G. Milburn,

J. Salzman, and H. Wiseman. CT is funded by a USA National Science Foundation Math and Physical Sciences Distinguished International Postdoctoral Research Fellowship. This work was supported by the Australian Research Council, the Australian Government and by the US National Security Agency (NSA), Advanced Research and Development Activity (ARDA) and the Army Research Office (ARO) under Contracts Nos. W911NF-04-1-0290 and W911NF-05-1-0284.

COMPETING FINANCIAL INTERESTS

The authors declare that they have no competing financial interests.

* Electronic address: andrew.greentree@ph.unimbelb.edu.au

[1] Jaynes E.T. & Cummings F.W., Comparison of Quantum and Semiclassical Radiation Theory with Application to the Beam Maser. *Proc. IEEE* **51**, 89-109 (1663).

[2] Birnbaum K.M., Boca A., Miller R., Boozer A.D., Northup T.E. & Kimble H.J., Photon blockade in an optical cavity with one trapped atom. *Nature* **436**, 87-90 (2005).

[3] Imamoğlu A., Schmidt H., Woods G. & Deutsch M., Strongly interacting photons in a nonlinear cavity. *Phys. Rev. Lett.* **79**, 1467-1470 (1997).

[4] Grangier P, Walls D & Gheri K, Comment on Strongly interacting photons in a nonlinear cavity. *Phys. Rev. Lett.* **81**, 2833 (1998).

[5] Greentree A.D., Vaccaro J.A., de Echaniz S.R. , Durrant A.V. & Marangos J.P., Prospects for photon blockade in four-level systems in the N configuration with more than one atom. *J. Opt. B: Quantum Semiclass. Opt.* **2**, 252259 (2000).

[6] Rebić S., Parkins A.S. & Tan S.M., Photon statistics of a single-atom intracavity system involving electromagnetically induced transparency. *Phys. Rev. A* **65**, 063804 (2002).

[7] Lang R.J. & Yariv A., An exact formulation of coupled-mode theory for coupled-cavity lasers. *IEEE J. Quantum Electron.* **24**, 66-72 (1988).

[8] Ozbay E., Bayindir M., Bulu I. & Cubukcu E., Investigation of localized coupled-cavity modes in two-dimensional photonic band gap structures. *IEEE J. Quantum Electron.* **38**, 837843 (2002).

[9] Greentree A.D., Salzman J., Prawer S. & Hollenberg L.C.L., Quantum gate for Q switching in monolithic photonic bandgap cavities containing two-level atoms, *Phys. Rev. A* **73**, 013818 (2006).

[10] Lee P.A. & Ramakrishnan T.V., Disordered electronic systems. *Rev. Mod. Phys.* **57**, 287-337 (1985).

[11] Fisher M.P.A., Weichman P.B., Grinstein G. & Fisher D.S., Boson localization and the superfluid-insulator transition. *Phys. Rev. B* **40**, 546-570 (1989).

[12] Jaksch D., Bruder C., Cirac J.I., Gardiner C.W. & Zoller P., Cold bosonic atoms in optical lattices. *Phys. Rev. Lett.* **81**, 3108 (1998).

[13] Cohen-Tannoudji C., Atoms in strong resonant fields. *Frontiers in Laser Spectroscopy*, eds. Balian R., Haroche S., & Liberman S., 1-104 (Elsevier Science Publishers B. V., North-Holland 1977).

[14] Hussin V. & Nieto L.M., Ladder operators and coherent states for the Jaynes-Cummings model in the rotating wave approximation, *Journal of Mathematical Physics* **46**, 122102 (2005).

[15] van Oosten D., van der Straten P. & Stoof H.T.C., Quantum phases in an optical lattice. *Phys. Rev. A* **63**, 053601 (2001).

[16] van Oosten D., van der Straten P. & Krishnamurthy H.R., Mott insulators in an optical lattice with high filling factors. *Phys. Rev. A* **67**, 033606 (2003).

[17] Sheshadri K., Krishnamurthy H.R., Pandit R. & Ramakrishnan T.V., Superfluid and insulating phases in an interacting-boson model: mean-field theory and the RPA. *Europhys. Lett.* **4**, 257 (1993).

[18] Krauth W. & Trivendi N., Mott and superfluid transitions in a strongly interacting lattice boson system. *Europhys. Lett.* **14**, 627 (1991).

[19] Krauth W., Trivedi N. & Ceperley D., Superfluid-insulator transition in disordered boson systems. *Phys. Rev. Lett.* **67**, 2307-2310 (1991).

[20] Xie Z.W. and Liu W.M., SuperfluidMott-insulator transition of dipolar bosons in an optical lattice. *Phys. Rev. A* **70**, 045602 (2004).

[21] Lewenstein M., Santos L., Baranov M.A. & Fehrmann H., Atomic Bose-Fermi mixtures in an optical lattice, *Phys. Rev. Lett.* **92**, 050401 (2004).

[22] Fehrmann H., Baranov M.A., Damski B., Lewenstein M. & Santos L., Mean-field theory of Bose-Fermi mixtures in optical lattices <Preprint at http://arXiv.org/abs/cond-mat/0307635> (2004).

[23] Littlewood P.B., Eastham P.R., Kneeling J.M.J., Marchetti F.M., Simons B.D. & Szymanska M.H., Models of coherent exciton condensation. *J. Phys. Condens. Matter* **16**, S3597-S3620 (2004).

[24] Olivero P. *et al.*, Ion beam assisted lift-off technique for three-dimensional micromachining of free standing single-crystal diamond. *Advanced Materials* **17**, 2427-2430 (2005).

[25] Baldwin J.W., Zalalutdinov M., Feygelson T., Butler J.E. & Houston B.H., Fabrication of short-wavelength photonic crystals in wide-band-gap nanocrystalline diamond films. *J. Vac. Sci. Technol. B* **24**, 50 (2006).

[26] Tomljenovic-Hanic S., Steel M.J., de Sterke C.M. & Salzman J., Diamond based photonic crystal microcavities. *Opt. Express* **14**, 3556 (2006).

[27] Greentree A.D. *et al.*, Critical components for diamond-based quantum coherent devices. *J. Phys. Condens. Matter* **18**, S825-S842 (2006).

[28] Song B.-S., Noda S., Asano T. & Akahane Y., Ultra-high-Q photonic double-heterostructure nanocavity, *Nat. Mater.* **4**, 207 (2005).

[29] Jamieson D.N. *et al.* Controlled shallow single-ion implantation in silicon using an active substrate for sub-20-keV ions. *Appl. Phys. Lett.* **86**, 202101 (2005).

[30] Tamarat Ph. *et al.*, Stark shift control of single optical centers in diamond. <Preprint at http://arXiv.org/abs/quant-ph/0607170> (2006).

[31] Wallraff A., Schuster D.I., Blais A., Frunzio L., Huang R.-S., Majer J., Kumar S., Girvin S.M. & Schoelkopf R.J., Strong coupling of a single photon to a super-

conducting qubit using circuit quantum electrodynamics. *Nature(London)* **431**, 162-167 (2004).

[32] Weidinger M., Varcoe B.T., Heerlein R. & Walther H., Trapping States in the Micromaser. *Phys. Rev. Lett.* **82**, 3795 (1999).

[33] Trupke M., Hinds E.A., Eriksson S., Curtis E.A., Moktadir Z., Kukharenka E. & Kraft M., Microfabricated high-finesse optical cavity with open access and small volume. *Appl. Phys. Lett.* **87**, 211106 (2005).

[34] Freedman M., Nayak C., Shtengel K., Non-Abelian topological phases in an extended Hubbard model. <Preprint at http://arXiv.org/abs/cond-mat/0309120> (2003).

[35] Kosterlitz J.M. & Thouless D.J., Ordering, metastability and phase transitions in two-dimensional systems. *J. Phys. C* **6**, 1181-1203 (1973).

[36] Scarola V.W. & Das Sarma S., Quantum Phases of the Extended Bose-Hubbard Hamiltonian: The Possibility of a Supersolid State of Cold Atoms in Optical Lattices, *Phys. Rev. Lett.* **95**, 033003 (2005).

[37] Hartmann M.J., Brandão F. G. S. L. & Plenio M.B., Strongly interacting polaritons in coupled arrays of cavities. <Preprint at http://arXiv.org/abs/quant-ph/0606097> (2006).

[38] Angelakis D.G., Santos M.F. & Bose S., Photon blockade induced Mott transitions and XY spin models in coupled cavity arrays. <Preprint at http://arXiv.org/abs/quant-ph/0606159> (2006).

[39] Rokhsar D.S. & Kotliar B.G., Gutzwiller projection for bosons. *Phys. Rev. B* **44**, 10328 (1991).

[40] Krauth W., Caffarel M. & Bouchaud J.-P., Gutzwiller wave function for a model of strongly interacting bosons. *Phys. Rev. B* **45**, 3137-3140 (1992).



FIG. 1: (a) Schematic showing a two-dimensional array of photonic bandgap cavities, with each cavity containing a single two-level atom (spheres). The photonic bandgap is described by a periodic perturbations of the dielectric medium, in this case represented by the regions drilled through a thin membrane, for example via focussed ion beam. The spacing between the holes will be of order λ/η where λ is the optical wavelength and η the refractive index. The membrane should be of order $\lambda/(2\eta)$ thick to confine light in the plane. Cavities are defined by defects in the periodic modulation, i.e. where a rod has *not* been drilled. By introducing a periodic array of defects in the photonic bandgap structure we can realise a lattice of photonic bandgap cavities, i.e. a photonic bandgap cavity superlattice. Photons can hop from one cavity to either of the three nearest neighbours with a frequency (photon hopping rate) κ, illustrated for some of the cavities by the double headed arrows. To effect photon-photon repulsion, we insert a single two-state system into each cavity, represented here by the red spheres. If the membrane were fabricated in diamond, the two-state systems could be ion implanted nitrogen-vacancy (NV) centres. In this lattice we have chosen for each cavity to have three nearest neighbours, so that the regime where photonic repulsion dominates over hopping is easier to achieve. (b) Schematic showing some of the pertinent energy scales within each atom-cavity system.



FIG. 2: Eigenspectrum for a single atom in a high Q cavity, as a function of the atom-cavity detuning, centered around 0, i.e. we plot $E - n\omega$. The eigenspectrum splits naturally into two branches, corresponding to the dressed states, $|+, n\rangle$ (upper branch) and $|-, n\rangle$ (lower branch). Note that the ground state $|g, 0\rangle$, naturally appears to be a member of each branch, introducing a significant departure from usual Bose-Hubbard style dynamics within each branch. Excluding $|g, 0\rangle$, the branches anti-cross at $\Delta = 0$, with the splitting increasing with increasing excitation number, n, given by the usual Rabi frequency, $\beta\sqrt{n}$.



FIG. 3: Boundaries between Mott lobes in the limit of low tunneling (small κ) as a function of μ and Δ. The two lowest stable regions correspond to $|g,0\rangle$ on the left, and $|-,1\rangle$ on the right. The boundary between these two domains follows the line $\mu = E_{|-,1\rangle}$. Higher order states can be seen in the central region, defined by the critical chemical potentials $\mu_c(n)$ and correspond to $|-,2\rangle$, $|-,3\rangle$ etc.



FIG. 4: Slices showing the superfluid order parameter, ψ, as a function of the photon hopping frequency, κ, and the chemical potential, μ, for (a) detuning, $\Delta = 0$, (b) $\Delta = -2$, and (c) $\Delta = 2$. The diagrams show Mott insulator lobes, indicated by the regions of $\psi = 0$, where the lowest have been shown. Dominating the left hand edge (where photonic repulsion dominates over hopping) is the Mott insulator phase (denoted MI), and the superfluid phase is found on the right hand edge (denoted SF). The white contour in (a) corresponds to the region where ψ becomes nonzero, delineating the quantum phase transition.



FIG. 5: Plateaus with constant density, ρ, indicating regions with a definite number state excitations, as a function of the chemical potential, μ, and photon hopping frequency, κ, for detuning, $\Delta = 0$. The first three plateaus are indicated, and the ground state configurations correspond to $|g, 0\rangle$, $|-, 1\rangle$ and $|-, 2\rangle$ for 0, 1, and 2 excitations (photons in this case) respectively, with the plateaus shrinking in size with increasing excitation number. Regions with varying ρ have coherent states as the ground state configuration.